arXiv:1509.08812v1 [math.RA] 29 Sep 2015

AN ISOMORPHISM LEMMA FOR GRADED RINGS

JASON BELL AND JAMES J. ZHANG

Abstract. Let A and B be two connected graded algebras finitely generated in degree one. If A is isomorphic to B as ungraded algebras, then they are also isomorphic to each other as graded algebras.

Introduction

The isomorphism problem has been studied by several researchers in recent years [BJ, CPWZ, Ga]. In this paper we prove a result which is useful for some of the isomorphism problems concerning graded algebras.

Throughout this paper, we let k be a base field, and all vector spaces, algebras, and morphisms are over k.

Theorem 0.1. *Let A and B be two connected graded algebras finitely generated in degree 1. If $A \cong B$ as ungraded algebras, then $A \cong B$ as graded algebras.*

Equivalently, we have

Corollary 0.2. *Suppose that an algebra A has two graded algebra decompositions*

$$A = \bigoplus_{i=0}^{\infty} A_i = \bigoplus_{i=0}^{\infty} B_i$$

such that

(1) $A_0 = B_0 = k$,
(2) *A is generated by A_1 (respectively, by B_1), and*
(3) *either A_1 or B_1 is finite dimensional over k.*

Then there is an algebra automorphism $\phi : A \to A$ such that $\phi(A_i) = B_i$ for all i.

Let $\mathrm{Aut}(A)$ (respectively, $\mathrm{Aut}_{gr}(A)$) be the group of algebra automorphisms (respectively, graded algebra automorphisms) of A. We have an immediate consequence.

Corollary 0.3. *Retain the hypotheses of Corollary* 0.2. *If $\mathrm{Aut}(A) = \mathrm{Aut}_{gr}(A)$, then $A_i = B_i$ for all i.*

As an application, we give the following solution of an isomorphism problem.

Let $\{p_{ij} \mid i < j\}$ be a set of invertible scalars in $k^\times := k \setminus \{0\}$. By convention, let $p_{ii} = 1$ and $p_{ji} = p_{ij}^{-1}$ if $i < j$. Recall that the skew polynomial ring $k_{p_{ij}}[x_i, \ldots, x_n]$ is generated by $x_1, \ldots, x_n$ and subject to the relations

$$x_j x_i = p_{ij} x_i x_j$$

2000 *Mathematics Subject Classification.* Primary 16W20, 16W50.
Key words and phrases. Isomorphism problem, skew polynomial ring.

for all i, j. An elementary change of generators means that we replace the ordered set $\{x_1, \ldots, x_n\}$ by $\{a_1 x_{\sigma(1)}, \ldots, a_n x_{\sigma(n)}\}$ where $a_i \in k^\times$ and σ is a permutation in S_n. The following result is viewed as a partial generalization of [Ga, Theorem 7.4].

Theorem 0.4. *Suppose that $p_{ij} \neq 1$ for all $i \neq j$. Let A be a graded ring $k_{p_{ij}}[x_1, \ldots, x_n]/M$ where M is an ideal in $k_{p_{ij}}[x_1, \ldots, x_n]_{\geq 3}$ and B be a graded ring $k_{p'_{ij}}[x_1, \ldots, x_m]/N$ where N is an ideal in $k_{p'_{ij}}[x_1, \ldots, x_m]_{\geq 2}$. If A is isomorphic to B as ungraded algebras, then $n = m$ and there is a permutation $\sigma \in S_n$ such that $p'_{ij} = p_{\sigma(i)\sigma(j)}$ for all i, j. Furthermore, after an elementary change of generators in A, $A = B$.*

1. Proof of Theorem 0.1

First we introduce an invariant of an algebra, which is similar to the Jacobson radical. Let dim denote the k-vector space dimension. Let s be a positive integer and define the Jacobson radical with tangent dimension s to be

$$J_s(A) = \bigcap \{I \mid I \subseteq A \text{ such that } \dim A/I = 1 \text{ and } \dim I/I^2 = s\}.$$

(In the case of an empty intersection, we take $J_s(A) = A$.) An ideal I of A with the property $\dim A/I = 1$, and $\dim I/I^2 = s$ is called a codimension 1 ideal of tangent dimension s. In general, given a sequence of non-negative integers $(s_i)_{i\geq 0}$, we can define

$$J_{(s_i)}(A) = \bigcap \{I \mid I \subseteq A, \dim I^i/I^{i+1} = s_i, \text{for all } i\}.$$

The following lemma is easy.

Lemma 1.1. *Let A be an algebra finitely generated over k.*

1. *Suppose A is generated by d elements. If $s > d$, then $J_s(A) = A$.*
2. *Suppose A is connected graded and $\dim A_{\geq 1}/(A_{\geq 1})^2 = d$ (this implies that A is generated by d elements). Then $J_d(A) \subseteq A_{\geq 1}$.*
3. *Suppose that k is infinite. Then $J_{s+1}(A[t]) = (J_s(A))[t]$.*

Proof. (1) Let $\{f_1, \ldots, f_d\}$ be a set of algebra generators of A. If I is an ideal of codimension 1, then I is generated by $\{x_i\}_{i=1}^d$ where $x_i := f_i - \alpha_i$ for some $\alpha_i \in k$. Then A is generated by $\{x_i\}_{i=1}^d$ as an algebra and I^2 is generated by $\{x_i x_j\}_{i,j=1}^d$. Hence $\dim I/I^2 \leq d$. Thus there is no ideal I of codimension 1 such that $\dim I/I^2 = s > d$. The assertion follows.

(2) Clear.

(3) Let I be an ideal of A of codimension 1 such that $\dim I/I^2 = s$. Let I_α be the ideal of $A[t]$ generated by I and $t - \alpha$. Then I_α is of codimension 1 such that $\dim I_\alpha/I_\alpha^2 = s + 1$. Using the fact that k is infinite, it is easy to check that

$$\bigcap_{\alpha \in k} I_\alpha = I[t]. \tag{E1.1.1}$$

Let J be any idea of $A[t]$ of codimension 1 such that $\dim J/J^2 = s + 1$. Let $I = J \cap A$. Then I is an ideal of A of codimension 1. There is always an $\alpha \in k$ such that $t - \alpha \in J$. Then J is generated by I and $t - \alpha$. So $J = I_\alpha$. Thus $\dim J/J^2 = \dim I/I^2 + 1$. So $\dim I/I^2 = s$. Now taking the intersection with all such I, equation (E1.1.1) implies that $J_{s+1}(A[t]) = J_s(A)[t]$. □

The next lemma is a special case of Theorem 0.1. We prove it first as a warm-up.

Lemma 1.2. *Let A and B be connected graded algebras generated in degree 1. Suppose that $d = \dim A_1 < \infty$ and $A_{\geq 1}$ is the unique codimension 1 ideal with tangent dimension d. If $\phi : A \to B$ is an isomorphism as ungraded algebras, then $A \cong B$ as graded algebras.*

Proof. By Lemma 1.1(1,2), $\dim A_1 = \dim B_1$. Since $A \cong B$, by hypothesis, there is a unique ideal of B of codimension 1 such that the tangent dimension is d. Therefore $B_{\geq 1}$ is the unique codimension 1 ideal of tangent dimension d. Thus ϕ maps $I := A_{\geq 1}$ to $L := B_{\geq 1}$. Therefore ϕ induces a graded algebra isomorphism from $\mathrm{gr}_I A := \bigoplus_{i=0}^{\infty} I^i/I^{j+1}$ to $\mathrm{gr}_L B := \bigoplus_{i=0}^{\infty} L^i/L^{j+1}$. Since $A \cong \mathrm{gr}_I A$ and $B \cong \mathrm{gr}_L B$ as graded algebras, there is a graded algebra isomorphism from A to B. □

Proof of Theorem 0.1. By Lemma 1.1(1,2), $\dim A_1 = \dim B_1 =: d < \infty$. Pick k-linear bases of A_1 and B_1 respectively, say, $\{x_1, \ldots, x_d\}$ and $\{y_1, \ldots, y_d\}$. Let $I = A_{\geq 1}$, $J = \phi(I)$, $L = B_{\geq 1}$, and $K = \phi^{-1}(L)$. Since $L = (y_1, \ldots, y_d)$ and B is connected graded, $L/L^2 = \bigoplus_{i=1}^{d} k\overline{y_i}$. Since K is a codimension 1 ideal of tangent dimension d, by making a linear change of the x_i, we may assume that $K = (x_1, \ldots, x_{d-1}, x_d - \alpha)$, for some $\alpha \in k$, and $K/K^2 = \bigoplus_{i=1}^{d-1} k\overline{x_i} \bigoplus k\overline{x_d - \alpha}$. The algebra isomorphism ϕ induces an k-linear isomorphism

$$\overline{\phi} : K/K^2 \to L/L^2.$$

By making a linear change of the y_i, we have that

$$\overline{\phi}(\overline{x_i}) = \overline{y_i}, \quad \text{for all } i = 1, \ldots, d-1, \quad \text{and} \quad \overline{\phi}(\overline{x_d - \alpha}) = \overline{y_d}.$$

Equivalently, we have that $\phi(x_i) = y_i + y_i'$ with $y_i' \in L^2$ for $i = 1, \ldots, d-1$ and that $\phi(x_d - \alpha) = y_d + y_d'$ with $y_d' \in L^2$.

The easy case is when $\alpha = 0$, then we have $J \subseteq L$. Since J has codimension 1, we then see that $J = L$. In fact, $\alpha = 0$ if and only if $J = L$, if and only if $K = I$. Since $\phi(I) = L$, the argument in the proof of Lemma 1.2 shows that

$$A \cong \mathrm{gr}_I(A) \cong \mathrm{gr}_L(B) \cong B.$$

The hard case is when α is nonzero (or equivalently, $J \neq L$ or $K \neq I$). Recall that $\phi(x_i) = y_i + y_i'$ with $y_i' \in L^2$ for $i = 1, \ldots, d-1$ and that $\phi(x_d) = \alpha + y_d + y_d'$ with $y_d' \in L^2$. For $i = 1, \ldots, d-1$ and $j \geq 1$, let $x_i^{[j]}$ denote the homogeneous element $[x_d, [x_d, \ldots, [x_d, x_i] \cdots]]$ where there are $j-1$ copies of x_d appeared in the expression. For example, $x_i^{[1]} = x_i$ for all $i = 1, \ldots, d-1$. Similarly we define $y_i^{[j]}$. Note that $x_i^{[j]}$ has degree j and $\phi(x_i^{[j]}) - y_i^{[j]} \in L^{j+1}$.

Let $r(x_i^{[j]})$ be a homogeneous relation of degree m in $x_i^{[j]}$ for $1 \leq i \leq d-1$, $j \geq 1$. Then applying ϕ, we see that $r(y_i^{[j]}) = r(y_i^{[j]}) - \phi(r(x_i^{[j]})) \in L^{m+1}$ and hence

$$r(y_i^{[j]}) = 0 \tag{E1.2.1}$$

in B.

In general, any homogeneous (noncommutative) polynomial $r := r(x_1, \ldots, x_d)$ in x_1, …, x_d of degree m can be written as

$$r = \sum_{s=0}^{m} x_d^s r_s(x_i^{[j]}) \tag{E1.2.2}$$

where $r_s(x_i^{[j]})$ is homogeneous polynomial of degree $m-s$ in $x_i^{[j]}$ for $1 \leq i \leq d-1$ and $j \geq 1$. Next we will prove the following two statements by induction:

Claim 1: If r is a homogeneous relation of degree n in $x_1, \ldots, x_d$ (so $r(x_1, \ldots, x_d) = 0$ in A), then $r(y_1, \ldots, y_d) = 0$ in B.

Claim 2: $\dim A_n = \dim B_n$.

In fact, we will use induction on n for all such isomorphisms $\phi : A \to B$ with the property that $\phi(I) \neq L$. There is nothing to prove for $n = 0$ and 1. Now assume that Claim 1 and Claim 2 hold for all $n < m$. To prove Claim 1 for $n = m$, suppose that we have a homogeneous relation r in $x_1, \ldots, x_d$ of degree m (so $r(x_1, \ldots, x_d) = 0$ in A) and we have a decomposition $r = \sum_{s=0}^{m} x_d^s r_s$ as described in (E1.2.2). We need show that $r(y_1, \ldots, y_d) = 0$.

Let t be the largest integer for which $r_t(x_i^{[j]})$ in (E1.2.2) is nonzero in A. (Equivalently the degree of $r_t(x_i^{[j]})$ is the smallest). If $t = 0$, then it follows that $r(x_1, \ldots, x_d)$ is a relation in $x_i^{[j]}$ But we have shown that all such relations have the property that $r(y_1, \ldots, y_d) = 0$, see (E1.2.1), as desired. Now we assume that $t > 0$. Applying ϕ to r, we see that

$$0 = \phi(r(x_1, \ldots, x_d)) = \sum_{s=0}^{t} (y_d + \alpha + y_d')^s \phi(r_s(x_i^{[j]})). \tag{E1.2.3}$$

Then, by construction and (E1.2.3), $\alpha^t r_t(y_i^{[j]}) \in L^{m-t+1}$. This then gives by homogeneity that $r_t(y_i^{[j]}) = 0$. We have just shown that there is some homogeneous relation r' in $y_1, \ldots, y_d$ of degree $m - t$ such that $r'(x_1, \ldots, x_d) \neq 0$. Since A_{m-t} and B_{m-t} have the same dimension by Claim 2 (for $n = m - t$), it follows that there must exist a relation r'' in $x_1, \ldots, x_d$ of degree $m - t$ such that $r''(y_1, \ldots, y_d)$ is nonzero. This contradicts Claim 1 for $n = m - t$. Therefore Claim 1 holds for $n = m$.

As a consequence of Claim 1 for $n = m$, $\dim A_m \geq \dim B_m$. Since Claim 2 is independent of the choices of bases $\{x_i\}_{i=1}^d$ and $\{y_i\}_{i=1}^d$, we obtain $\dim B_m \geq \dim A_m$ by applying the consequence to the map $\phi^{-1} : B \to A$. Therefore Claim 2 holds for $n = m$. This finishes the induction.

By Claims 1 and 2, since A is graded, the map $\phi : A \to B$ defined by $\phi(x_i) = y_i$ gives a graded algebra isomorphism from A to B as required. □

Proofs of Corollaries 0.2 and 0.3 are easy and omitted. Theorem 0.1 fails when A and B are not generated in degree 1.

Example 1.3. Let A be the algebra $k_{-1}[x_1, x_2] \otimes k[y_1, y_2]$ with $\deg x_1 = \deg x_2 = 1$ and $\deg y_1 = \deg y_2 = 2$. Let B be the algebra $k_{-1}[x_1, x_2] \otimes k[y_1, y_2]$ with $\deg x_1 = \deg x_2 = 2$ and $\deg y_1 = \deg y_2 = 1$. Then $A \cong B$ as ungraded algebras, but not as graded algebras. Further A and B have the same Hilbert series.

2. Application 1: isomorphism problem

In this section we consider the isomorphism problem for skew polynomial rings and their graded factor rings, and prove Theorem 0.4.

Lemma 2.1. *Let A be the skew polynomial ring $k_{p_{ij}}[x_1, \ldots, x_n]$ and let $B = A/I$ where I be a graded ideal of A contained in $A_{\geq 3}$. Suppose $p_{ij} \neq 1$ for all $i \neq j$.*

Then every normal element in B_1 is of the form cx_i for some scalar c and for some $1 \leq i \leq n$.

Proof. Let $f \in B_1 = A_1$. If $I = 0$, the assertion follows from [KKZ, Lemma 3.5(d)]. Since $I \subseteq A_{\geq 3}$, f is normal in A if and only if the image of f is normal in $B = A/I$. Therefore the assertion follows. □

Now we can prove Theorem 0.4.

Proof of Theorem 0.4. By Theorem 0.1, A is isomorphic to B as graded algebras. In particular, $n = m$. Let $\phi : B \to A$ be a graded algebra isomorphism. Since $\phi(x_i)$ are normal elements in A of degree 1, by Lemma 2.1, $\phi(x_i) = c_i x_{\sigma(i)}$ for some $c_i \in k$ and some permutation $\sigma \in S_n$. Up to an elementary change of basis of A, ϕ sends x_i to x_i for all i. The assertions follow. □

3. Application 2: cancellation problem

We have another quick application. Let $Z(A)$ be the center of an algebra A.

Theorem 3.1. *Let A and B be two connected graded algebras finitely generated in degree 1. Suppose that $Z(A) \cap A_1 = \{0\}$. If $A[t_1, \ldots, t_n] \cong B[s_1, \ldots, s_n]$ as ungraded algebras, then $A \cong B$.*

Proof. If we set $\deg t_i = \deg s_i = 1$ for all i, both $C := A[t_1, \ldots, t_n]$ and $D := B[s_1, \ldots, s_n]$ are connected graded and finitely generated in degree 1. Since $C \cong D$, by Theorem 0.1, there is a graded algebra isomorphism $\phi : C \cong D$. Since $Z(A) \cap A_1 = \{0\}$, $Z(C) \cap C_1 = \oplus_{i=1}^n k t_i$. By definition, the s_j are in the center of D. Therefore $\phi^{-1}(s_j) \in \oplus_{i=1}^n k t_i$ for all j. By a dimension argument, $\phi^{-1}(\oplus_{i=1}^n k s_i) = \oplus_{i=1}^n k t_i$. Modulo s_i and t_i we obtain an induced algebra isomorphism $\overline{\phi} : A \cong C/(t_i) \cong D/(s_i) \cong B$. □

Recall that an algebra A is *cancellative*, if, for any algebra B, an algebra isomorphism $A[t] \cong B[s]$ implies that $A \cong B$. Theorem 3.1 provides a weaker version of cancellation by assuming that both A and B are connected graded finitely generated in degree 1. It would be interesting if one can improve Theorem 3.1 by removing the hypothesis on B.

Acknowledgments

J. Bell was supported by NSERC Discovery Grant 326532-2011. J.J. Zhang was supported by the US National Science Foundation (grant Nos. DMS 0855743 and DMS 1402863).

References

[BJ] V.V. Bavula and D.A. Jordan, Isomorphism problems and groups of automorphisms for generalized Weyl algebras. *Trans. Amer. Math. Soc.* **353** (2001), no. 2, 769–794.

[CPWZ] S. Ceken, J. Palmieri, Y.-H. Wang and J.J. Zhang, Invariant theory for quantum Weyl algebras under finite group action, preprint, (2015), arXiv:1501.07881.

[Ga] J. Gaddis, Isomorphisms of some quantum spaces, *Ring theory and its applications*, 107–116, Contemp. Math., **609**, Amer. Math. Soc., Providence, RI, 2014.

[KKZ] E. Kirkman, J. Kuzmanovich and J.J. Zhang, Shephard-Todd-Chevalley Theorem for Skew Polynomial Rings, *Algebr. Represent. Theory* **13** (2010) no. 2, 127–158.

Bell: Department of Pure Mathematics, University of Waterloo, Waterloo, ON N2L 3G1, Canada
E-mail address: jpbell@uwaterloo.ca

Zhang: Department of Mathematics, Box 354350, University of Washington, Seattle, Washington 98195, USA
E-mail address: zhang@math.washington.edu